



BIAA Telerehabilitation Survey: Preliminary Results

Table 1. Discipline-Specific Clinical Data

Physical Therapy	Occupational Therapy	Speech/Language Therapy (Cognition)	Education Therapy	Counseling	Nursing
Timed Up and Go Test	Range of motion; Reaction time	Attention Process Training	Coin identification	Beck Depression Inventory	Medication counseling
6-Minute Walk Test; 10 Minute Walk Test	DASH; Fugl Meyer; Barthel & Modified Barthel Index	CVLT, Boston Naming Test, Selective Reminding Test	Adding change to \$1	Beck Anxiety Inventory	Tapering meds & maintaining behavior
Berg Balance Score	Quality of hand movements	Following Verbal Directions	SRA Reading	EMDR stages	Accuracy in medication ID
Functional Gait Assessment	Typing speed, coin flip test	RBANS, COWA, Woodcock-Johnson, Orientation Log	Computational Math	MMSE; MOCA; NeruoQol	Accuracy in filling pill boxes
Distance Walked	Canadian Occupational Performance Measure	Accuracy in articulation; Word Fluency	Glencoe Spelling Series	Sleep hygiene	Blood pressure monitoring
Tinetti Gait Score	Initiation & participation in ADL's	Dietary upgrades	Reading Fluency; Reading Comp	CAGE; Sobriety Strategies	Blood Glucose monitoring
Strength & Endurance data	Rapid Alternating Movements	Verbal/auditory memory; prospective memory	Listening skills; Critical Thinking	Social skills	Diet – low sodium, low carb, etc.